GENERALIZED RADON POTENTIAL

by Douglas Sprinkel
UTAH GEOLOGICAL AND MINERAL SU
Open-File Report 108 (revision of 9/12

GENERALIZED SOURCE OF RADON

Belt of rock types associated with above average concentrations of Uranit and susceptible of producing significant amounts of Radon. Belt configuration is patterns of Geologic Map of Utah (Hintze, 1980) and Uranium Equivalent Map of Utah (U. USGS). The Wasatch fault zone is also outlined as a generalized source of Radon

POINT SOURCE OF RADON

Localized source of enriched Uranium (U^{226}), or Radium (Ra^{226}) that occurs naturally or has been concentrated by human activities. These sources mostly occur as uranium mine dumps, uranium mill tailings, or deposits around thermal activity. Building on these point sources or using their deposits for building or backfill materials is unwise.

POINT SOURCES

- X Uranium Mines (active or inactive)
- ■ Uranium Mills (active or inactive)
- ▲ Selected Thermal Springs
- ▼ Geothermal Resource Areas

Snowville
Trenton
Richmond
Bear Lake
Rosette
Tremonton
Grouse Creek
Logan
Nibley
Randolph
Corinna
Paradise
Brigham City
Woodruff
Perry
Lucin
Great Salt Lake
Ogden
Plain City
Layton
Morgan
Coalville
Wendover
Salt Lake City
Delle
Kearns
Sandy
Park City
Tooele
Heber City
Vernal
American Fork
Roosevelt
Utah Lake
Provo
Dugway
Duchesne
Spanish Fork
Payson
Elberta
Thistle
Nephi
Trout Creek
Price
Levan
Mt. Pleasant
Delta
Woodside
Manti
Scipio
Castle Dale
Gunnison
Ferron
Sevier Lake
Fillmore
Green River
Cisco
Richfield
Moab
Koosharem
Marysvale
Loa
Milford
Hanksville
Beaver
Lund
Boulder
Monticello
Panguitch
Escalante
Newcastle
Cedar City
Tropic
Blanding
Hurricane
St. George
Mexican Hat
Kanab