Geology of the Burns Knoll Quadrangle, Beaver and Iron Counties, Utah

by Lehi F. Hintze, Myron G. Best, and Clark L. Weaver

1988

Quaternary mapping by Hintze, igneous rocks identified by Best, older bedrock mapping by Weaver (1980*)
*Brigham Young University Geology Studies, volume 27, part 3.

Plate 1
Utah Geological and Mineral Survey
Open-File Report 179
May 1990

This open-file release makes information available to the public which will not appear in another published form but is considered to be of value. It may not necessarily conform to formal UGS policy, technical review, or editorial standards, and therefore it may be premature for an individual or group to take action based on the contents of this report.

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

Map units

Qal	Alluvium
Qaf_1	Young fan deposits
Qaf_2	Middle fan deposits
Qaf_3	Old fan deposits
Qap	Pediment alluvium
Qla	Lacustrine and alluvial deposits
Qlu	Lacustrine deposits
Qeb	Eolian blowout
Qea	Eolian and alluvial deposits
Qe	Eolian deposits
QTa	Old alluvium
	UNCONFORMITY
Tbm	Blawn mafics
Ti	Isom Formation
Tsh	Andesite of Shauntie Hills
	UNCONFORMITY
Jt	Temple Cap Fm
Jn	Navajo Ss (quartzite)
Trc	Chinle Fm (hornfels)
Trs	Shinarump Cg
Trm	Moenkopi Fm (hornfels)
	THRUST
€s	Swasey Limestone
--B···	Bonneville shoreline

Qe/QTa indicates thin cover of Qe over QTa

Mapped, edited, and published by the Geological Survey

Control by USGS and USC&GS

Topography by photogrammetric methods from aerial photographs taken 1970. Field checked 1971

Projection and 10,000-foot grid ticks: Utah coordinate system, south zone (Lambert conformal conic) 1000-meter Universal Transverse Mercator grid ticks, zone 12, shown in blue. 1927 North American datum

Fine red dashed lines indicate selected fence lines

UTM GRID AND 1971 MAGNETIC NORTH DECLINATION AT CENTER OF SHEET

SCALE 1:24 000

1 0 1 MILE
1000 0 1000 2000 3000 4000 5000 6000 7000 FEET
1 .5 0 1 KILOMETER

CONTOUR INTERVAL 10 FEET
DOTTED LINES REPRESENT 5-FOOT CONTOURS
NATIONAL GEODETIC VERTICAL DATUM OF 1929

ROAD CLASSIFICATION


Primary highway, hard surface — Light-duty road, hard or improved surface
Secondary highway, hard surface — Unimproved road
Interstate Route — U. S. Route — State Route

UTAH
QUADRANGLE LOCATION

THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS
FOR SALE BY U. S. GEOLOGICAL SURVEY, DENVER, COLORADO 80225, OR RESTON, VIRGINIA 22092
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

Hintze + Best 1989
Burns Knoll quadrangle
1:24000

BURNS KNOLL, UTAH
N3807.5—W11315/7.5
1971
AMS 3360 III NE—SERIES V897



Sub-Quaternary rocks are probably mostly Tertiary igneous rocks, either mafic lavas or a pluton, as inferred from the prominent aeromagnetic high centered on the northern two-thirds of the quadrangle. Mesozoic and Paleozoic strata may also be present locally. Amount of vertical displacement on the Moonshine Well fault is not known but may be several thousand feet.