M. Rass January 22, 1993

Review of the Skinner Peaks Quadrangle Text and Geologic Map

TEXT

General comments -- There is too much redundance and wordiness throughout the text. Try to be concise to reduce the wordiness. Reorganization of the text will reduce the reduncance.

There are too many semicolons used throughout the text.

1. The abstract is poorly written. It is a general listing of stratigraphy, structure, surficial deposits, resources, and hazards. It also contains information on depositional environments of the map units that are from references. An abstract is not a listing and/or presentation of "reference" ideas. Therefore, it needs to be rewritten. The following reference will help you in writing a good abstract: Cochran, Wendell, 1979, <u>Geowriting:</u> a guide to writing, editing, and printing in earth science.

2. Ignore this number. My comment No. 1 supercedes what was originally discussed in this suggestion.

3. Intro, p.2; The manuscript needs a location figure with regional structures and pertinent location information included. Included in the Introduction are the following: geologic setting land use information, terrain description, accessibility info., date and length of the project, that it was part of a MS degree fullfillment, and the sponsoring university.

4. Stratigraphy, p.4; Move paragraph from p.24 to stratigraphy section intro. on p.4.

5. Arapien Shale, p.5-6; No mention or discussion of salt/evaporites/over-pressured shales in this section to enhance or support the later discussion of diapirism. Why is the Arapien diapiric?

6. North Horn Fm, p.7; rewrite and add sentence, Conglomerate is (clast-supported or matrix-supported?) and poorly sorted, with the gravel clasts in a poorly sorted, fine- to medium-grained calcareous sandstone. Gravel clasts are subangular to ....

7. Green River Fm, p.12; Use the terms <u>indurated</u>, <u>cemented</u> instead of incoherent when discussing the consolidation or cementation of a rock. Coherent is in the AGI Glossary but in my opinion is a poor choice.

8. Interpretation .., p.15; Having a schematic stratigraphic section as a figure would aid your discussion in this section.

9. p.26; Move your Qaf (Qaf3, of Clark) deposit discussion to the QUARTERNARY section under Alluvial Fan Deposits. I would

label this deposit Qafo and discuss it first under the Alluvial Fan Deposits section. Your Qacf2 would become Qaf and your Qacf1 would become Qafy.

10. p.29; Include all discussion of alluvial fan deposits in one subject heading.

Structure, p.32; Change the title of this section to 11. Structural Geology. In addition, I suggest the following rewrite to the intro. paragraph: The structural geology of the rocks in the area around the Skinner Peaks quadrangle is interpreted to be the result of a sequence of tectonic events beginning in the Cretaceous and continuing to the Holocene. From oldest to youngest these events are: Cretaceous to Early Tertiary Sevier Orogeny folding and thrusting, Middle to Late Tertiary development of the Colorado Plateau, and Late Tertiary normal faulting (references). Happening concurrently with these events was local diapirism of the Arapien Shale (reference). The younger tectonic events produced structural features that were superimposed on the older ones resulting in complex and often confusing structural relationships.

12. Structural Geology, p.33; This reviewer is a bit confused with this discussion. I suggest the following rewrite for the discussion marked in GREEN boxes.

In addition, the unconformity between the Arapien Shale and strata of the North Horn, Green River, and Goldens Ranch Formations may be related to Sevier deformation (reference ?). [Question -- Is the contact between the Arapien Shale and the other units depositional or structural, or both?]

Paleotopographic highs produced by uplift over structurally thickened sections may have resulted in erosion of older strata. Standlee (1985, ....) suggests that thrusting and folding indirectly may have caused the local Indianola structural highs observed by Weiss (1969) and Mattox (1986) in nearby areas.

13. Structural Geology, p.34; Need schematic map view and xsectional view figures to aid in this confusing discussion of the monocline.

14. Diapirism of the Arapien Shale, p.37-39; This section has several fatal organizational, discussion, and interpretation problems. The crux of the problems is distinguishing between diapiric structures (diapirs) and collapse structures (related to dissolution collapse of the diapir). The condensed stratigraphic section and unconformities at Skinner Peaks may indicate the presence of a diapir or diapiric structure. The graben at Flat Canyon may have been modified by dissolution collapse(?) of the faulted subsurface diapir. The term "diapiric collapse" that is used in the discussion is incorrect. <u>Diapiric</u> refers to growth of a diapir, <u>collapse</u> refers to distruction of a diapir. The terms do not go together. Another problem is the lack of discussion of what kind of diapirism is occurring with the Arapien. Is diapirism of the Arapien related to salt/evaporites or over-pressured shales? Previous workers suggest conflicting percentages of salt in the Arapien. If the Arapien contains a low percentage of salt, how can it behave diapirically?

The title of and discussion in this section should be rethought. Maybe two sections: "Structural features associated with diapirism of the Arapien Shale" and "Dissolution collapse structures associated with the Arapien Shale". Several things to consider when beginning the discussion are roughed out as follows:

Based on what I've read, several pieces of circumstantial evidence suggest the presence of a diapir at Skinner Peaks. Unusual fault pattern and chaotic arrangement of fault blocks, 2) the presence of highly-deformed evaporitic(?) Arapien Shale, 3) a complex stratigraphic/structural contact between the Arapien and the younger rocks, 4) a possible onlap configuration of a condensed late Cretaceous to Early Tertiary stratagraphic section on the Arapien, and 5) the indication of shallower than normal lacustrine facies rocks (oncolites, etc...) in strata of the Green River Fm at Skinner Peaks (However, this last one is a bit shakey). The combination of these observations suggest the area at Skinner Peaks may be a diapiric node of Arapien Shale. However, I'm not convinced that the Flat Canyon Graben may have been modified by dissolution collapse of that diapir. Your going to need more evidence than salty well water and some vague karst features.

15. Include and cite more references throughout the text. At several locations in the text these are indicated.

16. The Stratigraphy section of the text and the stratigraphy symbols do not match. Problems extist with the North Horn and Flagstaff formations and KTu map symbol at Skinner Peak. This reviewer suggests dropping the KTu designation. Go ahead and map the formations based on your picks in the measured section. The section " Interpretation of Stratigraphy at Skinner Peaks" should be eliminated with the majority of the discussion going to the North Horn Fm subsection. M. Ross Modified Outline of Skinner Peaks Quadrangle Manuscript

ABSTRACT INTRODUCTION STRATIGRAPHY Jurassic Arapien Shale **Cretaceous** - Tertiary (North Horn vs KT undivided) North Horn Formation elimiate KTy from map Tertiary be gutsey take a stand! **Flagstaff Formation** Colton Formation Green River Formation Golden Ranch Formation Unit I: I E these have to match with map symbols Unit H.: Z Unit MI.: 3 Unit IV.: 4 Unit X.: 5 Tertiary - Quaternary Older Alluvial Fan Deposits Pediment - Mantle Alluvium Quaternary Alluvial Fan Deposits < Qafo (Qafo = Qafz of Clark, 1987) Qaf Qafy Lake Bonneville Deposits Colluvium and Mass-Movement Deposits Alluvium **IGNEOUS ROCKS** Tertiary Hornblende Monzonite Porphyry STRUCTURAL GEOLOGY Sevier Orogeny Compressional Deformation Diapirism of the Arapien Shale -) more to the end it it makes for easier discussion Development of the Gunnison Plateau Basin and Range Extensional Deformation **GEOLOGIC HISTORY** ECONOMIC GEOLOGY Oil and Gas Sand and Gravel Gypsum **Tuffaceous Rock** Manganese WATER RESOURCES

1

GEOLOGIC HAZARDS Earthquake Mass Movements (or Landslides) Karst Development Water Salinity Problems ACKNOWLEDGEMENTS REFERENCES APPENDIX INTERIM GEOLOGIC MAP OF THE SKINNER PEAKS QUADRANGLE, JUAB AND SANPETE COUNTIES, UTAH By Tracey J. Felger Department of Geology University of Minnesota-Duluth

Ross review copy

1c) \$

#### ABSTRACT

The Skinner Peaks quadrangle is located in central Utah, just west of the leading edge of the Sevier fold-and-thrust belt, and in the transition zone between the Colorado Plateau and the *physiographic privinces* Basin and Range. The stratigraphy and structure of the rocks in the quadrangle reflect several tectonic events, including the Sevier *development* Orogeny, formation of the Colorado Plateau, and Basin and Range extension. Local diapiric movement of the Arapien Shale, which probably was initiated by these major tectonic events, further modified the structure and affected the stratigraphy.

Exposed bedrock units in the quadrangle include sedimentary, and pyroclastic, and intrusive rocks that range in age from Middle Jurassic to Late Oligocene. An unconformity separates Middle Jurassic marine strata of the Arapien Shale from the overlying Cretaceous-Tertiary strata. These Cretaceous-Tertiary strata include, in ascending stratigraphic order, the North Horn, Flagstaff, Colton, Green River, and Goldens Ranch Formations.

Strata of the North Horn, Flagstaff, and Colton Formations

1

represent the alluvial fan and plain, lacustrine, and fluvial were prevalent in environments conditions that dominated the Sevier foreland basin during the Late 🦟 western parts of the Formation Cretaceous and Early Eocene. Eocene Green River strata records the inundation of the basin by Lake Uinta, and the volcaniclastic Goldens and associated Delimentation ecords Ranch Formation is representative of the widespread volcanism that was in western (late EDcene - Olicocene occurring throughout Utah during Oligocene time. Two small igneous It hornblende monzonite porphyry intrucle Inrussic stratin. intrusions also were mapped as were unconsolidated surficial deposits nexting lacustrine, fluvial, colluvial, alluvial fan, and landslide deposits vironments are present ranging in age from Late Tertiary to Recent.



See

Major structures in the quadrangle are the Sage Valley Fault, the Western Juab Valley Fault Zone, the Wasatch Fault Zone, the West Gunnison Monocline, the Juab Valley Graben, and Flat Canyon Graben. Economic deposits include sand and gravel, gypsum, tuff,

carbonate rock, manganese, and water. Earthquakes, mass movements, karst development, and groundwater contamination are potential geologic hazards in the Skinner Peaks quadrangle.

Rewrite See

### INTRODUCTION

The Skinner Peaks 7.5 minute quadrangle is located approximately (mtrid) 100 miles south of Salt Lake City in Juab and Sanpete Counties, central Utah. The quadrangle extends from 39° 22' 30" to 39° 30' ( north latitude, and from 111° 52' 30" to 112° west longitude. It lies in the transition zone between the Colorado Plateau and Basin and Range Provinces, the Colorado Plateau Province is represented by the Gunnison Plateau, which terminates just east of Utah Highway 28. In

Introduction need a location figure (s) showing

physiographic & structural features.

# Need a sentence tiking Gunnison Platom to San Pitch Mtris (structure) (physiographic)

addition to the Gunnison Plateau, the Skinner Peaks quadrangle also includes the southern end of the West Hills, Mills Gap, the South Hills, and part of Juab Valley. Total relief in the quadrangle is approximately 1,700 feet; base elevation is 5,000 feet above sea level.

The first geologic map of the Skinner Peaks quadrangle was made by James W. Vogel of Ohio State University in 1957. Vogel mapped the geology at a scale of 1:31,680 on an imprecise planimetric base map constructed from aerial photos, no suitable topographic map of the area existed at that time. Witkind and others (1987) included the Skinner Peaks quadrangle as part of the Manti 30' x 60' quadrangle, although most of the geology that appears on the Manti Sheet was compiled from Vogel's original work.

Other early investigations of the structure and stratigraphy of central Utah were conducted by E. M. Spieker (1946, 1949) and his students from Ohio State University (e.g., Zeller, 1949; Muessig, 1951; Vogel, 1957). Faculty and students from Ohio State, Brigham Young, and Northern Illinois Universities have continued to expand and modify Spieker's earlier work.

# This is poorly constructed. Cite work that directly pretning to your gandongle. Be more specific on work is its contribution to your study area. STRATIGRAPHY

Sedimentary, pyroclastic, and igneous rocks ranging in age from Middle Jurassic to Late Oligocene are exposed in the Skinner Figure Fig

I reter to a strategraphic colum figure is the text.

and two igneous intrusions. Unconsolidated lacustrine, fluvial, colluvial, alluvial fan, and mass-movement sediments ranging in age from Late Tertiary to Recent were mapped in addition to the bedrock units. See A 1.14 section added here.

Precambrian and Paleozoic strata are not exposed as bedrock in the quadrangle, but they are exposed in the nearby Valley Mountains, Canyon Range, and southern Wasatch Mountains (Hintze, 1975), well data indicate these strata also underlie the study area (Standlee, 1982). Although Precambrian and Paleozoic strata are not exposed in the study area, clasts of Precambrian and Paleozoic strata are prevalent in the conglomerates of the North Horn, Flagstaff, Colton, Green River, and Goldens Ranch Formations, and in the various unconsolidated Tertiary-Quaternary deposits.

### JURASSIC

### Arapien Shale

The Arapien Shale, which was deposited in a narrow seaway during In the contrast of the quadrante Jurassic Callovian time, is exposed east of Utah Highway 28 along the west flank of the Gunnison Plateau. It underlies Skinner Peaks, and it also is exposed in and adjacent to Little Salt Creek Canyon.

The Arapien is composed of grayish-green, thinky-bedded micrite limestone, micrite, and calcareous siltstone; thinky-bedded, rippled, calcareous sandstone, and grayish-green or red calcareous mudstone with locally occurring pods of gypsum. These rock types are the Arapien state as defined by representative of units B and C of Hardy (1952). Thinly-bedded siltstone, shale, and rippled sandstone matching the description of unit C occurs in both the Little Salt Creek Canyon & and Skinner Peaks vicinity. These beds locally contain fossils (Z. M. Matching) identified as Ostrea sp., an observation that is congruent a with that of Zeller (1949, p.19), who noted the occurrence of Ostrea sp. in unit C sandstone in upper Little Salt Creek Canyon.

Index map of region

In outcrop the Arapien shale ... generally occurs as highly folded, contorted and faulted strata ... (Vogel, 1957, p. 32) that weathers to form steep, rugged, sparsely vegetated, gray hills. Most of the units within the Arapien weather into small chips or thin \* plates, ledges occur locally where more resistant sandstone or siltstone is present.

the word units (Stratigraphic) relationships between the Arapien and adjacent (ad this spot rocks units are (complex). The base of the formation is not exposed within or  $\mathcal{L}$ is contrising w Units used adjacent to the study area; however, data collected from drill-holes to describe contormance intern Arapien in SE Juab County indicate that the Arapien is underlain conformably Shale strata. under by the Twin Creek Limestone (Sprinkel, 1982). This relationship can why the term be observed in outcrop in the Mona quadrangle, 15 miles NE of the complex? Skinner Peaks quadrangle. In normal sequences the Arapien is overlain diapiric contacts conformably by the Twist Gulch Formation however, in the Skinner fmalt contact etc .... Peaks quadrangle, the Arapien is most commonly overlain unconformably Forlan by the Green River Formation. Locally, it is overlain unconformably by the North Horn Formation or the Goldens Ranch Formation. These unconformable relationships are best observed immediately south of Little Salt Creek Canyon and on the Skinner Peaks themselves.

Determination of an accurate thickness for the Arapien has been

Within the La section there is no mention / discussion of / satt/evaporites/over-pressured shales to support later discussion of diapirism.

> hampered by poor exposure (Sprinkel, 1982) and the intense deformation of the strata (Sprinkel, 1982; Standlee, 1982); estimates range from 3,000 to 11,000 feet throughout the area of its exposure (Eardley, 1933; Spieker, 1946; Hardy, 1952; Standlee; 1982). In this study, a thickness of approximately 440 feet was calculated from an incomplete, undeformed section of Arapien south of Little Salt Creek Canyon. Approximately 2,000 feet of Arapien was logged in a test hole in the NW corner of the quadrangle.

This discussion has serious problems of which I do not have the expertise to specifically comment. However my required knowledge allows me to recognize the overall problem. I suggest discussion it w/ PM (Gew).



### North Horn Formation

Large quantities of coarse-grained, clastic sediments were led from the Sevier Highland during the Late Cretaceous and Early Tertiary and deposited as a series of alluvial fans in the foreland basin to the east. These alluvial fans formed a conglomerate sequence that is represented by the Indianola Group, Price River Formation, and North Horn Formation. This sequence of conglomerates is almost 10,000 feet thick on the Gunnison Plateau (Hintze, 1988).

In the Skinner Peaks quadrangle, beds that tentatively have been identified as North Horn Formation are exposed in a narrow band on the NE side of Skinner Peaks. The North Horn Formation is not exposed anywhere else in the quadrangle, although it does crop out in the West Hills just north of the NW corner of the quadrangle (in the Juab (Inth(MST) durther the North than for the quadrangle). It also occurs in the subsurface in Juab Valley (Clark, ~ 1987).

Your discussion in the "Interpretation of the Stratigraphy of Skinner Peaks" would be better suited here ad contrined with this information bimictic - term not found in AGI glossory or sederal clastic sedemartology books use bimodal, polymodal?

Outcrops of North Horn Formation in the Skinner Peaks quadrangle are composed of poorly sorted, <u>bimictic</u> cliff- and ledge-forming # conglomerate. Clasts are subangular to subrounded pebbles, cobbles, # and boulders of purple and tan quartzite and dark blue-gray carbonate. Purple clasts were derived from the Precambrian Mutual Formation, and tan clasts were derived from the Cambrian Tintic Quartzite; dark blue-gray carbonates represent a variety of Paleozoic formations. Matrix is poorly-sorted, medium- to fine-grained, calcareous

sandstone. Clast size decreases up-section, the top of the section consists ( of interbedded conglomerate and sandstone. There is also an increase in the quartzite-to-carbonate clast ratio up-section, the lower part ( of the section has a 0%/100% carbonate/quartzite clast ratio, whereas the top of the section has a 75%/25% carbonate/quartzite clast ratio. The color of the unit also varies in an up-section direction; it is gray at the base, red in the middle, and gray at the top. The description of this section of North Horn is similar to Mattox's (1986, p. 80) description of "high escarpment and inner canyon" North Horn strata.

In most sections, especially farther east, the North Horn \* Formation lies conformably on top of the Price River Formation, and is in turn conformably overlain by the Flagstaff Formation; however, in the Skinner Peaks quadrangle, the North Horn Formation lies unconformably on top of the Jurassic Arapien Shale, and the relationship between it and the overlying strata is unclear.

The thickness of the North Horn Formation is also anomalous. The

exposed section on Skinner Peaks is only 300 feet thick however, only when KThh 6 miles to the west in the West Hills, Clark (1987) reported a thickness of approximately 800 feet, and approximately 1,700 feet of North Horn Formation was logged in a test hole just south of Chicken (reference) Creek Reservoir. No P The drastic thickness variations and the relationship between the North Horn Formation and adjacent units is discussed in detail in the

Section"Interpretation of the Stratigraphy of Skinner Peaks".

#### TERTIARY

## Flagstaff Formation phase of

The Flagstaff Formation represents a major lacustrine phase of deposition that occurred between the alluvial fan and floodplain deposition conditions represented by the North Horn Formation and the Colton Formation. Strata of the Flagstaff Formation range in age from Paleocene to Eocene; this age range is based primarily on paleontologic evidence that has been gathered by various workers throughout central Utah (LaRocque, 1951; Newman, 1974; Fouch and others, 1982).

In the Skinner Peaks quadrangle, the Flagstaff Formation is exposed in the east-dipping cuestas of the West Hills in the NW corner of the quadrangle. Beds tentatively identified as Flagstaff Formation also are exposed along the NE side of Skinner Peaks and are discussed in the "Interpretation of the Stratigraphy of Skinner Peaks".

A section of Flagstaff Formation was measured in the West Hills (onsists of (in during abundance) north of Mills Gap, Calcareous mudstone, sandstone, sandy limestone,

### (Felger, 19\_).

limestone, and conglomerate (listed in order of decreasing abundance) are the major rock types in this section. These strata are equivalent to the carbonate-clastic facies defined by Clark (1987) in the Juab quadrangle to the north.

The color of the strata varies from grayish-yellow to pale reddish-orange, with various hues of yellow being most common. The calcareous mudstone is massive; it weathers to a slope and ranges from 20-80 feet in thickness. The sandstone is usually calcareous and composed of medium- to coarse-grained quartz and lithic sand; locally, it is cross-bedded. Compositionally, the sandstones are quartz arenites, sublitharenites, and lithic arenites (Clark, 1987; Auby, proba 1985). Beds of sandstone form ledges that are 1-4 feet thick, and commonly are laterally discontinuous. Massive beds of sandy limestone and limestone form resistant ledges 2-20 feet thick; locally, these carbonate units are platy, weathering to slopes with local ledges. Beds of clast-supported conglomerate and conglomeratic sandstone occur locally throughout the section. These units are laterally discontinuous, often channel-form in shape, and 1-10 feet thick. \* Clasts are subangular to subrounded, poorly-sorted pebbles and cobbles of quartzite and sandstone. The matrix is medium- to coarse-grained calcareous sandstone that is composed of quartz and lithic sand.

The relative abundance of coarse-grained clastic material, the presence of cross-bedded sandstone, and the lateral discontinuity of the sandstone and conglomerate beds suggests that the Flagstaff Formation in the Mills Gap section was deposited in a near-shore, shallow-water environment. This interpretation is consistent with

allevial deposites?

9

those of Muessig (1951), Lambert (1976), and Clark (1987).

The base of the Flagstaff Formation is not exposed in the West Hills within the Skinner Peaks quadrangle; however, it is exposed in the Juab quadrangle to the north, and there the contact with the underlying North Horn is conformable and gradational (Clark, 1987), as is the contact between the Flagstaff and the overlying Colton Formation. The Flagstaff Formation is approximately 525 feet thick.

### Colton Formation

Fluvial and alluvial plain sediments, which are assigned to the Colton Formation, represent the final infilling of the Sevier foreland basin which occurred during the Early Eccene.

In the Skinner Peaks quadrangle, the Colton Formation is exposed in a conspicuous red swath in the east-dipping cuestas of the West Hills. Beds that tentatively have been identified in this study as Colton Formation are exposed on Skinner Peaks; and are discussed in LK to ET the "Interpretation of the Stratigraphy of Skinner Peaks".

In the West Hills in the Skinner Peaks quadrangle, the Colton e Formation is composed of reddish-brown mudstone, sandstone, and conglomerate; thin beds of limestone occur locally throughout the section and are considered to be the deposits of short-lived local poorly The Colton Formation as a whole is not well indurated, and it \*\* lakes. weathers to form a saddle between the more resistant Flagstaff Colton Formation see below Limestone) and Green River Formation. The mudstone is calcareous and -e weathers to a slope. The sandstone is friable and weathers to a slope 🥓 The sound stone is friable with locally occurring ledges. It is calcareous and is composed of Cotton Formation mudstone and soundstone is calcaveous and generally weathers to form a slope. with occassional ledges.

subrounded, medium- to coarse-grained quartz, feldspar, lithic fragments, and mica. Studies by Marcantel and Weiss (1968) and Stanley and Collinson (1979) show that Colton sandstones are commonly finer grained and contain greater amounts of mica and feldspar than the sandstones in the Flagstaff Formation. Beds of limestone are sandy, and they occur locally as low, discontinuous ledges.

binictic see p.7

The conglomerate (figure 1) is clast-supported, moderately sorted, and (bimictic) clasts are subrounded pebbles of approximately equal amounts of purple and tan quartzite (from the Mutual Formation and Tintic Quartzite), and dark blue-gray Paleozoic limestone. This Provenance suite of clasts indicates derivation from the Sevier Highland to the 🦀 west. The matrix, which comprises approximately 20 percent of the rock, is sandstone that is calcite-cemented and composed of medium- to L cemented by calcite coarse-grained, quartz and lithic sand. Conglomerate beds are 5 to 10 feet thick, channel-form, and laterally discontinuous; they occur as ledges and cliffs. Regionally, conglomerate is rare in the Colton and proximal to the source regions it occurs here only because the area was close to the edge of the along the western margin of the basin. basin.

The high percentage of mudstone, laterally discontinuous beds of conglomerate, sandstone, and limestone, and the red color of the strata attest to the fluvial (floodplain and channel) origin of the Colton Formation (Marcantel and Weiss, 1968).

In the West Hills in the Skinner Peaks quadrangle, the Colton Formation is underlain conformably by the Flagstaff Formation, and overlain conformably by the Green River Formation. The formation is approximately 300 feet thick.

11

Green River Formation 19

and marginal Sediments that were deposited in Lake Uinta from the Early \* make up through Late Eocene formed the strata of the Green River Formation. In the Skinner Peaks quadrangle, strata of the Green River Formation represent lake margin depositional environments. reflect the lake-marginal location of the quadrangle, and four distinct lithofacies are recognized/ from the base of the unit upward, they are the mudstone, clastic, and mudstone-micrite lithofacies of Clark (1987), and the Tawny facies of Zeller (1949). out croppings e The best exposures of strate of the mudstone, clastic, and mudstone-micrite lithofacies of the Green River Formation are in the cuestas of the West Hills, while the best exposures of the Tawny fith facies are found in the vicinity of Skinner Peaks. subbedding from your descussion Dee Eliminate Mudstone facies: The mudstone lithofacies is composed mostly of poorly indernites thinly bedded, grayish-yellow mudstone that is very incoherent and subsequently weathers to a slope. Thin, laterally discontinuous beds .eof quartzite pebble conglomerate and sandy limestone also occur locally throughout the unit. The unit is capped by a resistant bed of

The stromatolites occur as laterally-linked hemispheroids up to 2 feet in diameter.

stromatolitic limestone that contains brown and gray chert nodules.

<u>Clastic facies</u>: The clastic facies consists of conglomerate, conglomeratic sandstone, mudstone, and sandstone. The conglomerate and conglomeratic sandstone is reddish-brown or grayish-yellow/ it is <u>bimictic</u> with poorly-sorted pebbles and cobbles of quartzite and

e carbonate in a medium- to coarse-grained sand<del>stone</del> matrix. These conglomerate and conglomeratic sandstone units are poorly indurated and laterally discontinuous. Mudstones are reddish brown, thinly Lalcareous Sandstones are gray, calcite-cemented, and are laminated slope-formers. grains composed of quartz and lithic fragments; compositionally, these sandstones are sublitharenites, lithic arenites, and lithic wackes Sandstone beds form low ledges that are laterally (Clark, 1987). Beds of oolitic limestone that have been replaced by discontinuous. silica also occur locally throughout the clastic facies; ripple marks commonly are preserved on the tops of these oolitic beds.

Alternating beds of red or yellow mudstone, Mudstone-micrite facies: Imestone and yellow or gray micrite dominate the mudstone-micrite lithofacies. stThe mudstones are very thinly-bedded, poorly indurated, and, comprise greater than consequently, they weather to slopes) mudstones total over 50 percent ( of the mudstone-micrite facies (Clark, 1987). The micrite beds are well-indurated relatively coherent, and, consequently, they form a resistant cap over & the easily-eroded mudstones. These micrite beds are commonly platy d and fossiliferous; fossils include plant fragments, gastropods, and Clark (1987) noted pelecypods and ostracodes as well.

A thickness of 1,200 feet was calculated from outcrop width and bedding attitude for the Green River Formation in the West Hills of estimated This, thickness is approximately 300 the Skinner Peaks quadrangle. those feet greater than thicknesses calculated by Vogel (1957) and Clark The greater thuckness (1987) for the same general area. This suggests the presence of a found e fault in the section, but no evidence for a fault was seen in the

N

e

e

field.

sand vs

Sandstone

12 Tawny facies: Tawny (Beds consist of green, red, and variegated mudstone, and yellowish-tan coarse-grained sandstone, conglomerate, conglomeratic sandstone, and limestone. The sandstone is very cap Well indurated calcareous coherenty, it is usually cemented with calcite, and composed of quartz and minor amounts of lithic fragments. Sandstone beds form ledges that are several feet thick and laterally discontinuous / numerous (ap) vertebrate fossils are contained in sandstone beds near the top of the Channel-form beds of conglomerate and conglomeratic section. well indurated Clasts are subrounded to rounded sandstone also are very coherent. pebbles of dark blue-gray carbonate (>75%), and tan and purple quartzite (<25%); matrix is sandstone similar to that described above. Limestone is very dense and commonly fossiliferous, containing teeth and bone fragments, as well as gastropods of the species Australorbis (LaRocque, 1960). Strata of the Tawny facies match the description of strata in Millen's (1982) alluvial facies, which represents an alluvial or delta plain environment of deposition.

Fragments are generally used when describing angalor compounts at preceras, pyriclustic rocks, etc...

Granns is noundly used when describing subrounded clasts in a clastic rock.

Complex stratigraphic relationships separate the Tawny Beds from adjacent units. With the exception of Hunt (1950), all workers (Vogel, 1957; Millen, 1982; Norton, 1986) agree that the contact between the Tawny Beds and the underlying Green River Formation is conformable and gradational; this relationship was confirmed in this study as well. Tawny Beds also unconformably overlie the Arapien Shale south of Little Salt Creek Canyon. They are, in turn, overlain conformably by strata of the Goldens Ranch Formation.

14

This section is problematic. Most of this intermetion is redundat 15/ And section. This section & KTAh should be combraced. Some information (ie condensed stratigraphic section, unusually redunating features jete i. ) should be discussed in . Cha pirism Interpretation of the Stratigraphy of Skinner Peaks return different trom Stratigraphic sections in the Vicinity. Approximately 550 feet of conglomerate, thus, poorly understood. conglomeratic sandstone, sandstone, sandy limestone, and oncolitic limestone grade vertically into strata of the Tawny facies of the Green River Formation. Vogel (1957) and Witkind and others (1987) e mapped these strata as pad of the Tawny facies of the Green River more recent and rocks suggest they Formation. A closer evaluation of these units indicates that they a condensed section of to more accurately represent Late Cretaceous-Early Tertiary strata as (D.A. Spinkel, UGS, verbal communication, 19\_). suggested by Douglas A. Sprinkel of the Utah Ceological Survey (UGS). e 0 Evidence to support this interpretation is cited throughout the following section. Unit numbers (e.g., unit 4) correspond to the unit Measured numbers found in the Skinner Peaks Section in the Appendix. about 300 feet of A section of poorly sorted conglomerate and conglomeratic

sandstone, which is approximately 300 feet thick, lies unconformably on the Arapien Shale. These conglomerates were described in detail in the section on the North Horn Formation; only a summary description is presented here.

The conglomerate in the lower 220 feet of the section (unit 4) is massive, clast-supported, poorly-sorted, and bimictic. Clasts include subangular 10 subrounded pebbles, cobbles, and boulders of purple and tan quartzite, and a small percentage of dark blue-gray carbonate; matrix is poorly-sorted, medium- to fine-grained lithic sandstone. Clast size, and quartzite/carbonate clast ratio decreases up-section. The color of the unit also changes from gray to red up-section. This unit, which represents an alluvial fan deposit, is overlain by 55 feet

Figure w/ a schematic 15 Section would and this discussion. of interbedded conglomerate and sandstone (unit 5).

The conglomerate of unit 5 is gray, clast-supported, moderately-sorted, and bimictic. Clasts are subangular to subrounded cobbles of carbonate (75%) and quartzite (25%). The sandstone is composed of quartz; it is light-gray, medium-grained, well-sorted, and locally cross-bedded. This unit is indicative of an alluvial plain environment. Why is what advantant structures is texture, faces

The conglomerate sequence is overlain by approximately 100 feet of limestone (unit 6) and oncolitic limestone (unit 8; figure 2). The limestone is light-gray, massive, and finely-crystalline; it forms a ledge that is 10 feet thick. The oncolitic limestone, which contains oncolites up to three inches in diameter, forms cliffs and is 80 feet thick.

The oncolitic limestone is overlain by 110 feet of interbedded Sandy limestone and sandstone (unit 9) and interbedded sandstone and conglomerate (unit 10). The interbedded sandstone and sandy limestone is reddish-brown. The sandstone in this unit is calcareous and is composed of medium-grained quartz and minor amounts of lithic fragments; it forms local ledges throughout the slope-forming sandy limestone. This sequence is overlain by interbedded sandstone and conglomerate. The sandstone in this unit is also calcareous and is composed dominantly of medium-grained, well-sorted quartz sand. It also contains algal mat pieces and oncolites that may have been derived partially from the underlying oncolitic limestone. The conglomerate is clast-supported, moderately-sorted, and bimictic It is composed of approximately equal amounts of subrounded pebbles of

dark-blue-gray carbonate and purple and tan quartzite. Approximately 20 percent of the rock is matrix which is composed of quartz sandstone. Strata of these units represent a lake-marginal and fluvial environment which was typical of both the Flagstaff Formation and Colton Formation in this area; these strata grade vertically into the overlying Tawny Beds. The contacts between the lower units appear to be conformable.

Skinner Kenles A The section is a fining-upward sequence that represents a depositional transition through the following environments: alluvial fan (unit 4), alluvial plain (unit 5), lake-marginal and shallow-water lacustrine units exhibit lithologic and strufficialie (units 6-10). The lithology and stratigraphy of the units described above are characteristic of the North Horn, Flagstaff, and Colton Formations. It is difficult, however, to assign each unit to a specific formation. The conglomerates of units 4 and 5 match the regional description of North Horn strata. The limestone and oncolitic limestone of units 6-8 could be placed in either the North Horn Formation or the Flagstaff Formation. The sandy limestone, sandstone, and conglomerate of units 9 and 10 could be placed in either the Flagstaff Formation or Colton Formation, although the lack of a distinctive red color and abundant mudstone suggests that these Formation strata are more representative of the Flagstaff Limestone than they Formation \* il Regardless of which formation each unit is are of the Colton. assigned to, this section is far more representative of the regional sequence of Late Cretaceous-Early Tertiary strata than it is are the Towny Beds a formal name? representative of Tawny (Beds.

Based on this interpretation of the stratigraphy, very attenuated

10 this the best wird to use

why not include the Colton Fm?

sections of North Horn Formation and Flagstaff Formation are present on Skinner Peaks. The North Horn Formation is 300-400 feet thick placed depending on where the North Horn/Flagstaff contact is drawn. Likewise, the Flagstaff Formation is 110-220 feet thick. These thickness values are significantly less than values from the West Hills to the west and from the Gunnison Plateau to the east. The most logical explanation for the drastic thickness variations that occur over such a short distance is that welts of Arapien Shale formed local topographic highs in the basin during Late Cretaceous-Middle Tertiary This conclusion is supported by the presence of an unconformity time. between the Arapien Shale and Late Cretaceous-Early Tertiary strata and the presence of the oncolitic limestone. Oncolites, which are concretions of algae and sediment, form in shallow water, near-shore lacustrine environments. Weiss (1969) has shown that oncolites within the North Horn and Flagstaff Formations occur preferentially along what were actively-rising tectonic ridges.

Goldens Ranch Formation JC The onset of wide-spread volcanism in Utah occurred during the (Interme) Early Oligocenes This volcanism produced deposits, such as the A volcaniclastic Goldens Ranch Formation, which occurs throughout approximately one-third of the area of the Skinner Peaks quadrangle. In the western half of the quadrangle, the formation can be traced southward from the Chicken Creek Reservoir through the South Hills and In the hills on both the western and eastern sides of June Valley. 18 into the outcrops that flank the eastern side of the Sevier Bridge Reservoir. In the eastern half of the quadrangle, it occurs south of Chriss Canyon, and forms a "moat" that surrounds Skinner Peaks. Potassium-argon dates ranging from 38.5-29.9 m.y (Evernden and James, 1964; Witkind and Marvin, 1989) were obtained from samples collected from various units within the Chicken Creek Tuff Member. These dates confirm the Oligocene age of the formation.

en

e

For this report In the Skinner Peaks quadrangle, the Goldens Ranch Formation is within the quadrangle. Separated into five distinct, mappable units, (Units I-V, this study). In ascending order they are labeled Units 1-5. Units I through IV correspond to the Chicken Creek Tuff Member of Member? Meibos (1983), and unit V is the Hall Canyon Conglomerate or its of (reference) equivalent.

needed?

Unit I is an epiclastic conglomeratic sandstone (figure 3), ext. 4.4.4.4. The thickness of this unit is variable, ranging from 100 to approximately 500 feet thick. The contact between it and the underlying Eocene Green River Formation is gradational wherever it is exposed, as in the NE 1/4 of section 27, T. 16 S., R. 1 W.



Just above the contact with the Green River Formation, Unit I is composed of bentonitic shales interbedded with thin, platy limestone.

Confusio

Wirdunce

gradation

sandstone, forming a coarsening-upward sequence.

The upper three-quarters of Unit I are composed of sandstone and conglomeratic sandstone. The sandstone and matrix of the conglomeratic sandstone is most commonly a poorly-sorted lithic or arkosic sandstone. Grains are subangular, and range in size from 0.5-10 mm, with an average of 1 mm. The cement is typically calcareous, and the rock is friable to moderately coherent.

The coarsening-upward sequence of Unit I represents a shallow lacustrine/marginal lacustrine/fluvial environment of deposition that marks the end of Lake Uinta (De Vries and others, 1988).

2 varies from Unit II is a crystal vitric tuff that is 40-70 feet thick. Unit N: The contact between Unit I and Unit II is concordant and sharp. This e tuff is slightly welded, pink (weathered and fresh), and usually forms It is composed of 30-35 percent crystals and 65-70 percent slopes. pumice is not glassy glassy matrix. The crystals are euhedral and average 1 mm in size. Approximately 60 percent of these crystals are biotite, 40 percent are bipyramidal quartz, and sanidine occurs in trace amounts. The matrix ash (70-75%) and is composed of pumice fragments (25%-30%), which range in size from

0.5-20 mm, and ash (70%-75%). Bubble wall shards are visible in thin section.

relded ?

Unit NI: Unit III is coarse-grained epiclastic sandstone that is 50-90 feet thick. This unit is red or gray in color, forms resistant ledges and cliffs, and displays cross-bedding and channels. It is composed of approximately 60 percent bipyramidal quartz crystals, 5-15 . Ipercent lithic fragments, 15 percent sanidine, and traces of hematite. The lithic fragments are subrounded and range in size from 2-15 mm. The quartz crystals, hematite, and sanidine are subhedral to euhedral and average 2 mm in size. This unit is cemented by both silica and endersteel calcite, and is moderately to very coherent.

Unit II and Unit III are separated by an erosional contact. The detrifus nature of the contact and the presence of clasts of Unit II within Unit III suggest that Unit III was derived, at least in part, from the top of Unit II. Unit III represents a period of volcanic quiescence that occurred between the eruptive episodes that deposited Unit II and Unit IV.

Unit IV: Unit IV is an orange- or tan-colored vitric lithic tuff that is approximately 70-100 feet thick. The contact between it and Unit III is sharp and concordant. This tuff is less welded at the base where it weathers to form slopes; the upper part of the unit is better welded and it weathers to form vertical cliffs that commonly are cavernous.

The tuff of Unit IV is composed of 75 percent matrix, 20 percent

21

space

lithic fragments, and 5 percent crystals. The matrix is composed of 50 / percent ash and 50 percent pumice that ranges in size from 1-10 cm 2 and is commonly flattened in the bedding plane. The pumice forms a coarsening-upward sequence within the tuff. The lithic fragments are subangular to round, range in size from 0.5-2 cm and are composed of volcanic rocks and quartzite. Biotite, bipyramidal quartz, and a trace of sanidine constitute the crystal fraction of the tuff. These crystals are euhedral, and range in size from 0.5-2 mm.

5 a sandstone and complements sandstone sequence equivalent to the Unit V: Unit V is the Hall Canyon Conglomerate or its equivalent. It is is an epiclastic sandstone/conglomeratic sandstone of unknown thickness: In the Skinner Peaks quadrangle, the base of the unit is where its contact with Unit IV is crossical ad sharp. exposed in only one place, the topy is not exposed at all, due to of Unit V erosion, and the section is further, complicated by faulting. Clark (1987) reports that the thickness of the Hall Canyon Conglomerate to the jorth varies from 0-400 feet in the Juab quadrangle. The contact between Unit V and Unit IV is erosional and sharp.

The basal part of Unit V is an epiclastic sandstone that is very similar to Unit III; however, it is thin (rarely greater than 10 feet remainder thick), and contains sand-sized grains of Unit IV. The rest of Unit V & is very similar to Unit I in terms of texture and composition. The principal difference between Units I and V is the presence of angular tuffaccons clasts of Unit IV within Unit V. Unit V also contains more sandstone Unity exhibits. and less conglomeratic sandstone than Unit I. The sandstone is (medium-to conse-granned) (clast types). relatively homogeneous in terms of grain-size and composition (medium-The sandstone to coarse-grained lithic sandstone) it contains very large-scale,

tabular cross-bedding. The sedimentary structures, thickness, and overall stratigraphy of this unit suggest that it is an alluvial fan or a fan-delta deposit.

IGNEOUS ROCKS Hornblende Monzonite Porphyry Igneous Intrusions

Two small intrusions of hornblende monzonite porphyry occur in the Arapien Shale. One is located in the NW 1/4, NE 1/4 of section 36, T. 15 S., R. 1 W., and the other is located in the SW 1/4, SE 1/4 of section 25, T. 15 S., R. 1 W. These intrusions are not very resistant, and they weather to a grus-like talus that is black or dark-gray due to the abundance of hornblende. These and other intrusions in the vicinity were classified as dikes by Zeller (1949), Hunt (1950), and Vogel (1957). What do you think is their mighting

(Index of the fintrusions were examined under a

hornblende monzonite porpl webicates petrographic microscope. Approximately 65 percent of the rock is composed of phenocrysts, and the other 35 percent is a light-colored, 🦢 aphanitic groundmass of highly altered plagioclase and orthoclase. Approximately 75 percent of the phenocrysts are hornblende; feldspar and magnetite make up the remaining 59 percent. The hornblende cap phenocrysts occur as euhedral to subhedral laths that range from 0.01 plagioclase highly altered, yet Most feldspar phenocrysts are blocky, subhedral to 2.5 cm in length. crystals to euhedral, highly altered plagioclase crystals. Sallou

These intrusions are post-Jurassic in age based on the cross-cutting relationships in the Skinner Peaks quadrangle. Witkind and others (1987) cite an Oligocene(?) to Upper Eocene age for similar A santance on why they below the intrusions in the vicinity; however, the relationship of these Intrusions are Tertury.

23

Note to after STRATIGRAPHY section

intrusions to Tertiary units is not exposed in the Skinner Peaks quadrangle.

### TERTIARY-QUATERNARY

A variety of alluvial, colluvial, and lacustrine deposits blanket extensive areas of the Skinner Peaks quadrangle. These sediments Nove sectionents. range in age from Late Tertiary to Recent. They were deposited in response to tectonic and climatic events such as the development of the Gunnison Plateau and West Gunnison Monocline, the onset and continuation of Basin and Range faulting, and the advance and retreat of Lake Bonneville. Deposits (ATLF) -Mantle Quap Older Alluvial Fans and Pediment Alluvium Older detritus Sediment that was eroded from the Gunnison Plateau and West Gunnison Monocline was shed off to the west in a series of alluvial fans much like those that have formed in present-day Juab Valley. The 🥧 found in the southeaster uplifted remnants of the old alluvial fans are exposed along the flank of the quadragles of the West Gunnison Monocline in an area that extends from Broad Canyon to the southern end of the quadrangle. The material that forms yellowish-gray these deposits is semiconsolidated, massive to poorly-stratified, and poorly-sorted (ranging in size from sand to boulders), and Clasts are yellowish-gray in color. It is composed predominantly of sandstone, limestone, and conglomerate derived from the Green River Formation and includes clasts of pebbly sandstone from the Crazy Hollow Formation 4 and volcanic clasts derived from the Goldens Ranch Formation. The remnants of the old alluvial fans overlie the Goldens Ranch Formation, Green River Formation and Arapien Shale at various

24

Reverte into a complete sentence

where did this

Come from Z

elevations and reflect deposition over irregular paleptopography. This paleotopography may have been due in part to episodic Basin and Range faulting which began in the Miocene shortly after development of (reterence The thickness of these older alluvial fans the plateau and monocline. deposits varies from a few feet to 300 feet (Vogel, 1957). It is possible that these drastic thickness variations also reflect deposition over irregular paleotopography, with the thickest deposits representing paleo-lows and the thinner deposits representing paleo-highs. Pediment - Mantle Alluvin (QTap) -Martle Pediment alluvium, which caps the Goldens Ranch Formation in the Tapping an erosional surface developed on the Goldens Rouch South Hills, reflects an old erosional surface that developed during For may undicate an earlier period of uplift, erosion, and deposition in and after uplift of the South Hills area. The pediment alluvium, Varies from which is 0-20 feet thick, is very similar in texture and composition to the material that forms the old alluvial fans to the east. The most noticeable difference is the increased abundance of volcanic clasts and the local occurrence of red, semi- to moderatelyconsolidated, pebbly sandstone and sandy limestone. The red, pebbly sandstone and sandy limestone which occur locally as pods between the Goldens Ranch Formation and the poorly consolidated upper pediment alluvium may represent local ponds that formed on the erosional surface (Oviatt, personal communication, 1989). Like the older DIGE alluvial fans, the pediment alluvium occurs at relatively higher above the current drainage. elevations, reflecting the uplift and dissection that occurred after deposition.

ING

The distribution of the pediment alluvium and the alluvial fans reflects Lustig's (1969) prediction that areas with larger highlands favor alluvial fan development, and areas with lower highlands favor

25

### pediment development.

The age of the older alluvial fans and the pediment alluvium is \* Unknown and their correlation tentative. not known for certain. They are no older than Early Miocene because \* they formed after the development of the plateau and the onset of Basin and Range faulting. They are no younger than Earliest Pleistocene because Lake Bonneville sediments locally surround the me capped by these older

-maritle

e

bases of hills that these old alluvial deposits cap.

A solitary alluvial fan (mapped as (Qaf) in this study)

corresponding to Qaf<sub>3</sub> of Clark (1987) was mapped in the NW corner of by a younger fam and is faulted. the quadrangle. This fan is very dissected, faulted, and higher in The Ught-Brown aboved deform elevation than a younger fan which surrounds it. It is composed of light-brown, poorly-sorted, clay- to boulder-size material that is Many of the larger clasts are carbonete rock fribably derived from the Hagstoff fam (lake) subangular to subrounded. The poorly-sorted nature of the deposit, plus its proximity to the mouth of a deeply incised canyon that cuts through the Flagstaff Formation, indicate that this fan is a debris flow as Clark (1987) suggested. Clark (1987) estimates that the fan is at least 50 feet thick. Based on its relatively high elevation and on the very dissected and faulted nature of the fan, it formed either in the Latest Tertiary or Earliest Quaternary.

### it should be Labeled Qato

### QUATERNARY

Older Coalescing Alluvial Fans Deposits (Qaf)

Areas covered by old alluvial fans and pediment alluvium were differentially uplifted by Basin and Range faulting and then eroded, leaving only remnants of these old alluvial deposits capping the hills  $p_{e}t_{e}$ :  $t_{e}s$ along the flank of the monocline and in the South Hills. The material Muching uplifts is and that was eroded from these uplifted areas was deposited as a series of -(Raf and Rafy) coalescing alluvial fans that fill present-day Juab Valley. Material that was derived from the South and West Hills was shed primarily to the east, although some was deposited in the low spots to the west of the South Hills. Material derived from the Gunnison Plateau was shed into Juab Valley to the west. As Clark (1987) noted, the fans from the Gunnison Plateau are significantly larger than those emanating from the West and South Hills; consequently, the convergence line of the two fan systems lies west of the center of Juab Valley. Allword fan alluvium is reddish-brown to yellowish-gray,

unconsolidated, poorly-sorted, and massive to crudely bedded; local channels suggest a fluvial environment of deposition. Material is clay- to boulder-size, although sand- and pebble-size material is most As expected common; grain size decreases in a down-fan direction. Quartzite, limestone, sandstone, and volcanic rocks form the majority of the pebble- and cobble-size clasts. Data from a gravity survey (Zoback, 1983) across northern Juab Valley indicates that alluvial fan deposits are approximately 3,900 feet thick in that portion of the valley. *(reference)* Since Juab valley shallows to the south, the equivalent deposits in the Skinner Peaks area to the south are probably thinner than those to ~

The youngest sediment contained in the coalescing fans was detrifue A deposited on the fan surfaces during recent time; the oldest sediment of the back of the deposite, may have the oldest sediment of contained in these fans was probably deposited in the Late Tertiary, although there is no observable evidence to confirm this. Lake Bonneville sediments overlap coalescing fan deposits in the southwest

deposits

Qat

corner of the quadrangle, indicating that the deposits must be at older than late Pleistocene (latest Quaternary) least as old as Earliest Pleistocene.

Lake Bonneville Sediments

During the high stand of Lake Bonneville, which occurred 15,000 yrbp approximately 16,000-17,000 years ago, water from the lake spilled through Leamington Canyon, drowning the Sevier River and forming a fresh-water estuary (Oviatt, personal communication, 1989) that extended almost as far south as Redmond (Currey, 1982). The eastern shore of this estuary cut across the southwestern corner of the Skinner Peaks quadrangle. Sediments deposited in the estuary are exposed in the low, gently-sloping, dissected, fan-shaped patches in the Washboard and in wave-cut cliffs along the Sevier Bridge Reservoir. These sediments occur up to an elevation of 5,090 feet, which was the overflow elevation of the lake during the Bonneville Stage (Currey, 1982). A change in vegetation pattern that is best observed on aerial photos also occurs between 5,090-5,100 feet. It is presumed, based on this elevation, that this change in vegetation marks the shoreline of Lake Bonneville. It also is presumed, on the basis of elevation, that water from Lake Bonneville spilled through Mills Gap and flooded the Chicken Creek Reservoir area. / There are no deposits or shoreline features to substantiate this, but it is possible that Lake Bonneville sediments and shoreline features were there once but have been obliterated since by present-day Chicken Creek Reservoir. of bacustrine sediments

Although exposures are poor except along the Sevier Bridge

Reservoir, the sediments are fairly distinctive (especially on aerial photos) and can be distinguished from the surrounding alluvium, without much difficulty. Poor exposures obscure the nature of the contact between the Lake Bonneville sediments and the surrounding alluvium, but at one location (section 30, T. 16 S., R. 1 W.), the lake sediments clearly overlap the Quaternary-Tertiary pediment alluvium. Elsewhere (e.g., on the Washboard), the Bonneville sediments are slightly higher than the adjacent alluvium which suggests deposition of the Lake Bonneville sediments on top of the adjacent alluvium. This observation is consistent with the relationships observed by Mattox (1986) in the Hells Kitchen Canyon SE quadrangle, 10 miles southeast of the present study area.

The Bonneville sediments are light brown, unconsolidated, coarseto fine-grained sand, silt, and mud. These sediments form a fining-upward sequence that is 30-60 feet thick and are composed mostly of silt and mud. Deposits are finely laminated and cross-laminated; soft-sediment deformation structures and ripple cross-lamination are common near the base of the exposed section. These characteristics, combined with the lack of foreset and bottomset beds, fit Oviatt's (1984) description of underflow fan deposits, which are similar to deltaic deposits.

Younger Coalescing Alluvial Fan (Qafy)

in that they have coalesed into a fam

complex, however

A series of younger coalescing alluvial fans, rests on top of electronic older coalescing alluvial fans north of Little Salt Creek Canyon. The younger fans are very similar to their older counterparts, however,

Allavent tar

they are considerably smaller in size, and they slope more steeply toward the valley. The composition of these younger fans is also in that different from their older counterparts, most of the material is angular, pebble-size fragments of limestone that were derived from the Arapien Shale. These deposits are only 50-100 feet thick.

Younger alluvial fans, such as those that are found north of  $A = I_{o}$ Little Salt Creek Canyon, form in response to climatic or tectonic changes that lower base level (Pazzaglia and Wells, 1989; Bull, 1990). In the Skinner Peaks area, base level could have been lowered by the retreat of Lake Bonneville, continued Basin and Range faulting, or a combination of both of these events.

The very local occurrence of the younger alluvial fans suggests that they formed in response to renewed uplift along a fault segment and not in response to the regional lowering of base level that would have resulted from the retreat of Lake Bonneville. This hypothesis is supported by the presence of Recent fault scarps that cut the older coalescing alluvial fans; however, the older coalescing alluvial fans in Juab Valley and the Lake Bonneville sediments are incised by gullies that are as much as 15 feet deep, which suggests a regional lowering of base level. Perhaps the deep gullies are an expression of 105 ADDAS a regional lowering of base level that was due to the retreat of Lake recent movement Bonneville, and the younger alluvial fans reflect Recent Basin and along Range activity on a local fault segment. Assuming that these younger alluvial fans are related to the Basin and Range faulting that produced the fault scarps, the age of these fans is Late Pleistocene Are the fans younger than take Bonneville Migh stand ? to Recent.

Colluvium, Alluvium, and Landslide Deposits The youngest sediments in the quadrangle are colluvium, alluvium, and landslide deposits which are all Recent in age. The colluvium (Qc) forms steeply-sloping, cone-shaped deposits along the base of the slopes from which it was derived. It is unconsolidated, very angular, very poorly-sorted, clay to boulder size material. The color and composition of these deposits reflect the formation or formations from which they were derived. These deposits are 0-15 feet thick.

Mass-Movedent

The alluvium occurs along most drainages, at higher elevations, such as Flat Canyon and the South Hills, it forms broad, even surfaces of low relief. Like the colluvium, the composition and color of the alluvium reflect the local bedrock from which it was derived. In most cases, it is unconsolidated, gray or brown, in color and massive to poorly stratified. Alluvial material is clay- to cobble-size, subangular to subrounded, and poorly- to well-sorted. These deposits are generally less than 30 feet thick.

Two landslides are the only mass-movement deposits that were observed in the Skinner Peaks quadrangle. One of the landslides during located occurred on the north side of Chriss Canyon in the SE 1/4 of section 11, T. 16 S., R. 1 W., the other is located south of Skinner Peaks in the SE 1/4 of section 22, T. 16 S., R. 1 W. Both of these landslides occurred in strata of the Green River Formation and consequently are composed of very angular, poorly-sorted blocks of carbonate and sandstone in a matrix of mudstone. The Chriss Canyon landslide occurred in 1984 (Weiss, personal communication, 1989) after a period of heavy rain. Presumably the Skinner Peaks landslide, which is as

unsure of massive, is it = to featureless?
fresh as the Chriss Canyon landslide, also occurred in 1984.

STRUCTURE GEOLOGY interpreted to be rocks in the The structural geology of the Skinner Peaks quadrangle is the result of Sevier thrusting, formation of the Colorado Plateau, Basin (reterences) and Range faulting, and local diapirism of the Arapien Shale. The al features structures that were produced during one tectonic event were superimposed on the structures that formed during the previous tectonic event. This resulted in complex and confusing geologic relationships.

#### Sevier Thrusting

The Sevier Orogeny, which began in the Late Jurassic and continued into the Paleocene (Armstrong, 1968), was the first tectonic recognized event that affected the Skinner Peaks quadrangle. It was tolding and characterized by eastward-directed thrusting which placed Precambrian, upper Paleozoic, and lower Mesozoic strata over strata as young as Middle Jurassic. Middle Jurassic marine shales such as the Arapien are structurally incompetent and consequently acted as glide planes for the thrusting that built the Sevier Highland.

There is very little surface evidence of Sevier thrusting in the Skinner Peaks quadrangle; however, substantial subsurface evidence (Standlee, 1982; Lawton, 1985; Clark, 1987) indicates that some surface features can be attributed to the event. Data collected from isugglat drill-holes in and adjacent to the study area reveal several interpreted to stratigraphic repetitions. These repetitions indicate thrust faults John are correct. He

1.xart

32

owere

that formed during Sevier thrusting (Standlee, 1982; Lawton, 1985). Drastic variations of the thickness of the Arapien Shale and adjacent units are also attributed to thrusting.

The only surface evidence that can be attributed directly to Sevier thrusting is the highly contorted strata of the Arapien Shale. It is possible, however, that the unconformity that occurs between the Arapien Shale and strata of the North Horn, Green River, and Goldens Ranch Formations may be related to the Sevier orogenic event.

A recent study by Sims and Morris (1989) indicates that thrusting of a competent unit over an incompetent unit (e.g., the Sevier fold-and-thrust belt) will cause the incompetent unit to shorten and thicken close to the hinterland, and uplift will occur over the thickened region. As a result, the incompetent unit should be highly deformed, as is the Arapien Shale. Another possible result of this process is the formation of topographic highs in the area of

thickening. Standlee (1985, personal communication to S. Mattox) suggested that thrusting and folding indirectly may have caused the local Indianola highs observed by Weiss (1969) and Mattox (1986).

Vewrite

It is also possible that the paleo-highs are the result of diapiric movement of the Arapien Shale. Differential loading or tectonic activity is often necessary to initiate diapirism (Lemon, 1985; Jackson and Talbot, 1986); the influx of coarse-grained clastic material from the highland to the west and the eastward directed thrusting that was occurring at this time would have provided both of these mechanisms. The presence of a thick section of oncolitic limestone on Skinner Peaks supports the theory that this area was

why? explain actively rising during deposition. whether fectionic determition or dispirison or some combition

Regardless of which explanation is correct, it is certainly reasonable to conclude that the unconformity that occurs between the Arapien Shale and strata of the North Horn, Green River, and Goldens Ranch Formations is related to Sevier thrusting.

> at Whit Gunnison Morror Devie ONMEN Formation of the Gunnison Plateau

#### West Gunnison Monocline

There is not In the Skinner Peaks quadrangle, the Colorado Plateau Province is (San Pitch Mountains) whose western margin is represented by the Gunnison Plateau which terminates as the West Just Gunnison Monocline inside the east edge of the quadrangle. The West trends North-South and The Gunnison Monocline is approximately 18 miles long, and it extends from Fayette Wash in the Hells Kitchen Canyon SE quadrangle to Buck Canyon, Idmu for at the conference. (Mattox, 1986). NEED FIGURE At the surface In the Skinner Peaks quadrangle, the West Gunnison Monocline Westward consists of Green River Formation and Goldens Ranch Formationsstrata which dips 25 to 30 degrees to the west or southwest. Dips of 55 degrees and greater were observed in Green River strata on Skinner Peaks, but these values are anomalously high and may reflect diapiric modification by the underlying Arapien Shale.

> A thick section of Arapien Shale cores the monocline and extends eastward under the synclinal structure of the plateau. In general, the Arapien is highly deformed, and attitudes are quite variable. Attitudes measured in a relatively undeformed section below the Arapien-Green River unconformity south of Little Salt Creek Canyon dip consistently 40 to 45 degrees SE/ these attitudes are consistent with

France ( SMURG

those observed by Zeller (1949) in Arapien strata east of the Skinner Peaks quadrangle.

Based on the interpretations of Standlee (1982) and Lawton (1985), the Arapien core of the monocline represents a ramp structure that formed during Sevier thrusting; it is likely that the variable attitudes of the Arapien strata reflect deformation due to the thrusting event, as well as later modification by tectonically activated diapirism.

The West Gunnison Monocline and the Gunnison Plateau formed during Late Oligocene or Early Miocene time. The timing of this event & is constrained by the Oligocene Goldens Ranch Formation, which is represents the youngest strata on the monocline. The conformable contact between the Green River Formation and overlying Goldens Ranch Formation indicates that monoclinal warping had not begun prior to deposition of the Goldens Ranch Formation.

1 Mana

# Basin and Range Extension

The structural geology of the Skinner Peaks quadrangle is Aumerous dominated by north-south trending, high-angle normal faults, including MIC sufficient faults are: the Sage Valley Fault, the Western Juab Valley Fault Zone (WJVFZ), and the Wasatch Fault Zone (WFZ). Smaller normal faults also dissect the area. Why are these Bigh faults? Summerice into and make note of reformed, on Why they are Bigh faults?

Sage Valley Fault

The Sage Valley Fault is a high-angle, down-to-the-west fault which bounds the west side of the West Hills and the east side of Sage

Norma

### Within the NW corner of the quadrangle

Valley. The fault trends approximately N 10 E: Clark (1987) states that the fault has at least 2,900 feet of throw. Triangular facets that have formed along the western side of the West Hills define the fault scarp. The fault does not cut any Quaternary units within the Skinner Peaks quadrangle.

#### Western Juab Valley Fault Zone

The Western Juab Valley Fault Zone (WJVFZ) bounds the West Hills on the east and Juab Valley on the west. This fault is thought to be part of a zone of concealed down-to-the-east, high-angle normal faults. Surface evidence for the WJVFZ is sparse. Southeast of Chicken Creek Reservoir the fault appears to place upper Goldens Ranch Formation against Green River Formation and lower Goldens Ranch Formation. The fault which trends roughly N 40° E, has an estimated we throw of 1,000 feet.

#### Wasatch Fault Zone

The Wasatch Fault Zone (WFZ) bounds the west edge of the West Gunnison monocline and the east edge of Juab Valley. It is a zone of high-angle normal fault and is characterized by down-to-the-west movement. Triangular facets or faceted spurs of Arapien Shale south of Little Salt Creek Canyon Fault and fault scarps in Pleistocene alluvial fans attest to the presence of the fault. The fault scarps, which can be seen just west of Skinner Peaks, show approximately 5 to 10 feet of displacement. The Wasatch Fault trends approximately N 20° E and has an estimated throw of approximately 5,000 feet. Recent gravity and seismic data presented by Zoback (1983) indicate that Juab Valley, which is bounded on the west by the Western -Juab Valley Fault Zone and on the east by the Wasatch Fault Zone, is indicate that contains up to 3,000 feet of alluvial fill.

Other Faults

lena

Other faults that occur throughout the quadrangle include high-angle cross-faults such as those in the West Hills and the fault which parallels Old Botham Road in the South Hills area. These structures are possibly related to local strain accommodation that occurred during Basin and Range extension.

Other Structures

Basin and Range normal faulting not only produced the structures described above, it also affected the structure of the West Gunnison Monocline by dissecting the west-dipping strata into a series of west-dipping fault-blocks that are bounded by north-south-trending normal faults. Strata in the southern end of the quadrangle have been affected most noticeably.

Vertical joints, which trend approximately 30 degrees west and east of north, are prevalent in Green River and Goldens Ranch strata. The joints probably represent shear fractures that formed due to east-west extension.

Structural Diapirism of the Arapien Shale Evidence throughout the quadrangle indicates that diapiric movement of the Arapien Shale modified the structure of the area biocally. This local, episodic diapirism was probably initiated by tectonic events such as Sevier thrusting, development of the West Gunnison Monocline, and Basin and Range extension.

Sevier compression

Flat Canyon Graben and Skinner Peaks

Flat Canyon Graben is a structure that may represent an of an underly extensional graben that has been modified by diapiric collapse. This The graben is about I mile wide a diaDIV . structure is approximately one mile wide. It begins near Timber Canyon in the Hells Kitchen Canyon SE guadrangle and extends north to intering to the Stemmer Penk gradragle. Chriss Creek where it bends to the west. This graben is bounded on the east by the high-angle, down-to-the-west normal fault which parallels the southwest front of the Gunnison Plateau. It places Hall Canyon Conglomerate against Flagstaff and Green River strata. The west edge of the graben is bounded by a down-to-the-east normal fault which places the Hall Canyon Conglomerate against Green River and Arapien strata. outcrops and faults a, Westward The bend in the graben parallels the northwest trend of Skinner "These NW trends Peaks, which cuts across the otherwise north-south trending structures blocks that are related to the Basin and Range-Colorado Plateau provinces. characterIsTIC of Flat Cangon The graben, like Skinner Peaks, is underlain by Arapien Shale. The

The graben, like Skinner Peaks, is underlain by Arapien Shale. The presence of the Arapien in the subsurface beneath the Flat Canyon graben is manifest in salty well water and sink holes (W. Jay Dalley, landowner, personal communication, 1989). It seems reasonable to assume from this evidence that the structure of the Flat Canyon Graben and the adjacent Skinner Peaks is controlled in part by diapiric

why not is karst related to the Flagstatt or Tgr. collapse of the Arapien. It also seems reasonable to assume, based on the timing of the event, that the mobility of the Arapien was triggered by Basin and Range faulting.

#### Other Diapir Related Structures

Rootless fault blocks of Green River formation can be observed "floating" in Arapien Shale on the flanks of Skinner Peaks in the NE 1/4 of section 22 and the SW 1/4 of section 15 T. 15 S., R. 1 W. fault These blocks are similar to the detached blocks of Colton and Green River Formation described by Willis (1986) approximately 30 miles to the south in the Salina quadrangle. I concur with Willis' (1986) interpretation that these detached blocks are slump blocks which, in this case, slid off of the Skinner Peaks block. Is this a surfact process?

A small syncline in Green River strata that unconformably overlie the Arapien Shale in the NE corner of the Skinner Peaks quadrangle is also thought to have formed by diapiric movement of the Arapien O.A. (Sprinkel, personal communication, 1989). Contacts between the Arapien and overlying units are often sheared, with slickensides and well-foliated clays similar to those described by Willis (1986) in the Salina quadrangle. These contacts are also indicative of movement. Mapping Shuke.

#### ECONOMIC GEOLOGY

Economic deposits in the Skinner Peaks quadrangle and vicinity include sand and gravel, gypsum, tuff, carbonate rock, manganese, and petroleum products, and water. The sand and gravel occurs as pluid here alluvial, colluvial, and lacustrine deposits. Material ranges in size

road ballast (reference

incepte from clay to boulders; most material is sand and gravel composed of quartzite and carbonate clasts, with local concentrations of volcanic clasts. The sand and gravel, which is used primarily as road ballast, up is quarried from numerous gravel pits throughout the quadrangle.

MOVE

the useable size

Active quarrying of gypsum from the Arapien Shale on the NE side of Skinner Peaks began in 1989. This gypsum can be used in the production of dry-wall or as a bonding agent in cement.

Tuff from Unit IV ((Tvg,)) of the Goldens Ranch Formation formerly was guarried south of Skinner Peaks and in the Painted Rocks area for use as poultry grits, and soil mineralizer and conditioner (Vogel, This operation was run by the Azome Utah Mining Company of 1957). Sterling, Utah, and the products were marketed under the trade name "Azomite" (Vogel, 1957).

Carbonate rock that is found in the Flagstaff Limestone and Green River Formation possibly could be used as building or dimension stone. Unfortunately, in the Skinner Peaks quadrangle, neither of these formations contain sufficient amounts of limestone or dolomite to make quarrying a profitable economic venture because both formations contain anomalously high amounts of coarse-grained clastic material.

Small amounts of manganese occur in fault zones within the volcaniclastic Goldens Ranch Formation. The manganese occurs as dendritic pyrolusite in a calcite matrix. Pyrolusite is a secondary mineral that results from the alteration of manganese minerals (Edwards and Atkinson, 1986), which are present in small amounts in most crystalline rocks (Hurlbut and Klein, 1971). The manganese that ap forms the pyrolusite was probably leached from the surrounding Goldens

Ranch Formation and deposited with calcite along the fault zones.

Oil and gas exploration has taken place throughout central Utah because of the structural similarities between it and the producing and of the overthrust belt of Wyoming (Clark, 1987). Several oil companies have drilled test wells in Juab Valley and on the Gunnison Plateau in SE Juab County, no productive reservoirs have been discovered to date.

#### WATER RESOURCES

Water resources are somewhat limited in the Skinner Peaks quadrangle. Surface water occurs in the Chicken Creek and Sevier # Bridge Reservoirs, in Chicken Creek, and as small springs in the vicinity of the Skinner Peaks. Depth to the top of the water table is @ more than 100 feet (Bjorklund and Robinson, 1968) in the area of Juab Valley that lies between the South Hills and the west margin of the *for the formal and the formal for the formal for the formal for the formal for the formal formal for the formal formal* 

## GEOLOGIC HAZARDS - superate topics into subheadings

Earthquakes, mass movements, karst development, and groundwater contamination are the potential geologic hazards in the Skinner Peaks quadrangle and vicinity.

The Skinner Peaks quadrangle is centered roughly on the Wasatch Fault Zone which is part of the Intermountain seismic belt (McKee and Arabasz, 1982); the potential for catastrophic earthquakes is high. Gree Earthquakes may result in destructive ground shaking, surface rupture (le of alluvium, soil liquefaction, and differential settling (Clark, that effect empression of the structures 1987); they also may trigger mass movements such as snow avalanches Earthquakes and landslides. Landslides also may occur simply because strata are The Green River, etc., formations are prone to mass movement incompetent or poorly consolidated. Heavy rain or large volumes of whenomenon melt-water moving over steep, sparsely-vegetated mudstone slopes may of surface rocks result in mass wasting.

1/20 t

The development of karst topography and contamination of fresh groundwater are both related to the Arapien Shale. The evaporite-rich Arapien underlies much of the Skinner Peaks guadrangle. Groundwater moving through the Arapien dissolves the evaporates causing surface collapse and subsequent formation of sink-holes; evaporite dissolution also results in the contamination of the groundwater. Land-owner W. Jay Dailey reported the development of sink-holes and collapse structures in hay fields in Flat Canyon; he also reported salty water in a stock well in Flat Canyon. Vogel (1957) and Hunt (1950) cite similar reports from local residents concerning the quality of well More to after the STRUCTURAL GEOLOGY section

#### GEOLOGIC HISTORY AND INTERPRETATIONS

Aspects of the geologic history of the Skinner Peaks quadrangle were discussed throughout the stratigraphy and structural geology The following discussion is a sections of this manuscript. A brief synopsis of the geological and summary of history is presented here along with interpretations concerning the al geology structure and stratigraphy of the quadrangle.

geologic history of West-central Utah The Precambrian through Early Jurassic interval was dominated by shallow peralle deposits along enstern margin of the deposition of marine and continental sediments in the Cordilleran reterence. a. the surface These rocks are not exposed as bedrock in the miogeocline. are present quadrangle, but they do occur in the subsurface and as clasts in

conglomerates of the North Horn, Flagstaff, Colton, Green River, and hedrock Goldens Ranch Formations. The oldest exposed strata are the marine shales of the Middle Jurassic Arapien Shale. The sediments that l N-S trending comprise these strata were deposited by a shallow arm of the sea which 2 advanced from Canada, through central Utah, and into northern Arizona. (reterence) By the Late Jurassic this sea had retreated to the north. Compression A detormetion western margin of the caused by the subduction of the Pacific Plate under the North American reterence) Plate also started to affect central Utah around this time. migrated from west to east across western North American folding and Eastward-directed thrusting placed Precambrian, Paleozoic, and reaching Western Utabi by the Cheataceous. Mesozoic strata over the incompetent Arapien Shale which acted as a compressional deformation produced progenic This thrusting built the Sevier Highland and glide plane. Along its enstern edge corresponding foreland basin. reterence ). Durine

In Middle and Late Cretaceous time, the Skinner Peaks quadrangle, which was located in the foreland basin just east of the Sevier coarse-gramed detritus Highland, began to receive sediment that was being eroded from the highland and deposited in the basin as alluvial fans. Continued eastward sediment of the basin floor thrusting to the east and the differential loading that was caused by the influx of sediment from the west initiated diapiric movement of the evaporite-rich Arapien Shale. This local, episodic diapirism cap produced local topographic highs of Arapien Shale within the basin. Consequently, unconformities developed between the Arapien and various Cretaceous-Tertiary units that were being deposited in the foreland Based on the stratigraphic relationships and the abundance of basin. oncolitic limestone on Skinner Peaks, this area was the site of an actively rising topographic high of Arapien Shale.

The unconformity between the Arapien and the Green River

SK.

Formation indicates that tectonically activated diapirism continued into through the Early Tertiary, during which time the foreland basin was dominated by alternating lacustrine and fluvial conditions which produced the strata of the Flagstaff, Colton, and Green River formations. In the Skinner Peaks quadrangle, these formations have an anomalously high clastic fraction because the quadrangle was located along the western margin of the basin. Wide-spread volcanism dominated the landscape of central Utah in

the Oligocene, producing formations such as the volcaniclastic Goldens and the since t

Formation.

The Gunnison Plateau and the West Gunnison Monocline formed in Mrobably in the late the Late Oligocene after deposition of the Goldens Ranch Formation. Objourn Sediment was eroded from the plateau and monocline and deposited into

coalescing alluvial fans in the basin to the west.

No PNo P Basin and Range extension began shortly after the formation of produced mumerous the monocline. The extension dissected the area with north-south trending normal faults, such as the Sage Valley and Wasatch faults and produced east- and west-dipping fault blocks. Uplifted areas were dissected and eroded, and the sediment was deposited as alluvial fans to the adjacet grabus, such as in present-day Juab Valley.

In the Pleistocene, Lake Bonneville reached the Bonneville Stage, and (acastrice in the Skinner fick flooding the Sevier River and depositing underflow fan sediments. The About Approximately 2,000 years later the lake retreated catastrophically, lowering the regional base level. Active down-cutting through the

### pre-existing sufficial sedements

alluvial fans in Juab Valley and in stream gullies attests to the change in base level; continued Basin and Range extension also steepened the average regional gradient. Fault scarps that cut alluvial fan deposits, and the formation of secondary alluvial fans are evidence of Recent Basin and Range faulting. Total rewrite

#### ACKNOWLEDGEMENTS

This project, which was funded by the Utah Geological Survey, could not have been completed without the help of many people. The following people spent time in the field with me, edited my manuscript, or discussed my work with me: Dr. Timothy B. Holst, Dr. Richard W. Ojakangas, Dr. Wanda J. Taylor, and Nancy S. Nelson (University of Minnesota-Duluth); Dr. Malcolm P. Weiss, Steven R. Mattox, Dr. James A. Walker, and Rimmer De Vries (Northern Illinois University); Douglas A. Sprinkel, Michael L. Ross, Michael Schubat, & Grant C. Willis, and Lehi F. Hintze (Utah Geological Survey); Martin L. Sorensen, and Hal T. Morris (U.S. Geological Survey); and Dr. C.G. Oviatt (Kansas State University). The time that these people contributed was invaluable.

#### REFERENCES

Armstrong, R.L., 1968, Sevier orogenic belt in Nevada and Utah: Geological Society of America Bulletin, v. 79, p. 203-232.

Auby, W.L., 1985, Petrologic study of a measured section of Flagstaff Formation: Unpublished report, Northern Illinois University.

Bjorklund, L.J., and Robinson, G.B., 1968, Ground-water resources of the Sevier River Basin between Yuba Dam and Leamington Canyon, Utah: U.S. Geological Survey Water-Supply Paper 1848, 79 p.

Bull, W.B., 1990, Tectonic and climatic alluvial fans: Geological Society of America Abstracts with Programs, v. 22, no. 3, p. 11.

Clark, D.L., 1987, Geology of the Juab quadrangle, Juab County, Utah:

Unpublished M.S. thesis, Northern Illinois University, 323 p.

Currey, D.R., 1982, Lake Bonneville: Selected features of relevance to neotectonic analysis: United States Geological Survey Open-File Report 82-1070, 30 p.

Eardley, A.J., 1933, Stratigraphy of the southern Wasatch Mountains, Utah: Michigan Acad. Sci. Papers, v. 19, p. 377-400.

Edwards, R., and Atkinson, K., 1986, Ore Deposit Geology, University Press, Cambridge, 466 p.

Evernden, J.F., and James, G.T., 1964, Potassium-argon dates and the Tertiary floras of North America: American Journal of Science, v. 262, p. 945-971.

Fouch, T.D., Lawdon, T.F., Nichols, D.J., Cashion, W.B., and Cobban, W.A., 1982, Chart showing preliminary correlation of major Albian to Middle Eocene rock units from the Sanpete Valley in central Utah to the Book Cliffs in eastern Utah, in Nielson, D.L., editor, Overthrust belt of Utah: Utah Geological Association Publication 10, p. 267-272.

Hardy, C.T., 1952, Eastern Sevier Valley, Sevier and Sanpete counties, Utah with reference to formations of Jurassic age: Utah

Geological and Mineral Survey Bulletin 43, 98 p. or use Hinte, 1980 shte geologie map

Hintze, L.F., 1975, Geological highway map of Utah@ Brigham Young University Geology Studies, Special Publication 3, 1 sheet.
Hintze, L.F., 1988, Geologic history of Utah: Brigham Young University

Geology Studies, Special Publication 7, 202 p.

Hunt, R.E., 1950, The geology of the northern pad of the Gunnison Plateau, Utah: Unpublished Ph.D. dissertation, Ohio State University, 267 p.

Hurlbut, C.S., and Klein, C., 1977, Manual of Mineralogy, 19th edition (after J.D. Dana), John Wiley and Sons, New York, 532 p.

- Jackson, M.P.A., and Talbot, C.J., 1986, External strain rates and dynamics of salt structures: Geological Society of America Bulletin, v. 97, p. 305-323.
- Lambert, D.L., 1976, A detailed stratigraphic study of initial deposition of Tertiary lacustrine sediments near Mills, Utah: Brigham Young University Geology Studies, v. 23, pt. 3, p. 9-35.
- LaRocque, A.L., 1951, Molluscan fauna of the Flagstaff Formation, central Utah: Geological Society of America Bulletin, v. 62, p. 1457-1458.

LaRocque, A.L., 1960, Molluscan faunas of the Flagstaff Formation of central Utah: Geological Society of America Memoir 78, 100 p. Lawton, T.F., 1985, Style and timing of frontal structures, thrust belt, central Utah: American Association of Petroleum Geologists Bulletin, v. 69, no. 7, p. 1145-1159.

- Lemon, N.M., 1985, Physical modeling of sedimentation adjacent to diapirs and comparison with Late Precambrian Oratunga breccia body in central Flinders Ranges, South Australia: American Association of Petroleum Geologists Bulletin, v. 69, p. 1327-1338.
- Lustig, L.K., 1969, Trend surface analysis of the Basin and Range Province and some geomorphic implications: U.S. Geological Survey Professional Paper, 500D.
- Marcantel, E.L., and Weiss, M.P., 1968, Colton Formation (Eocene: fluviatile) and associated lacustrine beds, Gunnison Plateau, central Utah: Ohio Journal of Science, v. 68, no. 1, p. 40-49.
- Mattox, S.R., 1986, The geology of the Hells Kitchen Canyon SE quadrangle, Sanpete County, Utah: Unpublished M.S. thesis, Northern Illinois University, 448 p.
- McKee, M.E., and Arabasz, W.J., 1982, Microearthquake studies-across the Basin and Range-Colorado Plateau transition in central Utah, <u>in Nielson, D.L., editor, Overthrust belt of Utah: Utah</u> Geological Association Publication 10, p. 137-150.
- Meibos, L.C., 1983, Structure and stratigraphy of the Nephi NW 7 1/2-minute quadrangle, Juab County, Utah: Brigham Young University Geological Studies, v. 30, pt. 1, p. 37-58.
- Millen, T.M., 1982, Stratigraphy and petrology of the Green RiverFormation (Eocene), Gunnison Plateau, central Utah: UnpublishedM. S. thesis, Northern Illinois University, 220 p.

Muessig, S.J., 1951, Geology of a pad of Long Ridge, Utah: Unpublished Ph.D. dissertation, Ohio State University, 213 p.

- Newman, K.R., 1974, Palynomorph zones in early Tertiary formations of the Piceance Creek and Uinta basins, Colorado and Utah, Guidebook to the energy resources of the Piceance Creek basin, Colorado: Rocky Mountain Association of Geologists Guidebook 25th Annual Field Conference, p. 47-55.
- Norton, K.L., 1986, Paleogeography and lithofacies study of the Crazy Hollow Formation, central Utah: Unpublished M.S. thesis, Northern Illinois University, 183 p.
- Oviatt, C.G., 1984, Lake Bonneville stratigraphy at the Old River Bed and Leamington, Utah: Unpublished Ph.D. dissertation, University of Utah, 122 p.
- Pazzaglia, F.J., and Wells, S.G., 1989, Relative role of tectonism and climatic change on the evolution of Quaternary depositional landforms along a segmented range-front fault, Rio Grande rift: Geological Society of America Abstracts with Programs, v. 21, no. 6, p. A269.
- Sims, D., and Morris, A., 1989, Structural control in fold-thrust belts by low-competence units: Geological Society of America Abstracts with Programs, v. 21, no. 6, p. A135.
- Spieker, E.M., 1946, Late Mesozoic and early Cenozoic history of central Utah: U.S. Geological Suwey Professional Paper 205-D, p. 117-161.
- Spieker, E.M., 1949, The transition between the Colorado Plateau and the Great Basin in central Utah: Utah Geological Society

Guidebook 4, 106 p.

- Sprinkel, D.A., 1982, Twin Creek Limestone-Arapien Shale relations in central Utah, <u>in</u> Nielson, D.L., Overthrust belt of Utah: Utah Geological Association Publication 10, p. 169-180.
- Standlee, L.A., 1982, Structure and stratigraphy of Jurassic rocks in central Utah: their influence on tectonic development of the Cordilleran foreland thrust belt, <u>in</u> Power, R.B., editor, Geological Studies of the Cordilleran Thrust Belt: Rocky Mountain Association of Geologists, p. 357-382.
- Stanley, K.O., and Collinson, J.W., 1979, Depositional history of Paleocene-Lower Eocene Flagstaff Limestone and coeval rocks, central Utah: American Association of Petroleum Geologists Bulletin, v. 63, p. 357-382.
- Vogel, J.W., 1957, The geology of southernmost Juab Valley and adjacent highlands, Juab County, Utah: Unpublished M.S. thesis, Ohio State University, 152 p.
- Weiss, M.P., 1969, Oncolites, paleoecology, and Laramide tectonics, central Utah: American Association of Petroleum Geologists Bulletin, v. 53, p. 1105-1120.
- Willis, G.C., 1986, Geologic map of the Salina quadrangle, Sevier County, Utah: Utah Geological and Mineral Survey Map 83, 1:24,000.
- Witkind, I.J., and Marvin, R.F., 1989, Significance of new potassium-argon ages from the Goldens Ranch and Moroni Formations, Sanpete-Sevier Valley area, central Utah: Geological Society of America Bulletin, v. 101, p. 534-548.

- Witkind, I.J., Weiss, M.P., and Brown, T.L., 1987, Geologic map of the Manti 30 x 60 minute quadrangle, Carbon, Emery, Juab, Sanpete and Sevier counties, Utah: U.S. Geological Survey Miscellaneous Investigations Series Map 1-1631.
- Zeller, H.D., 1949, The geology of the west-central portion of the Gunnison Plateau: Unpublished M.S. thesis, Ohio State University, 83 p.
- Zoback, M.L., 1983, Structure and Cenozoic tectonism along the Wasatch fault zone, Utah, <u>in</u> Miller, D.M., and others, editors, Tectonic and stratigraphic studies in the eastern Great Basin region: Geological Society of America Memoir 157, p. 3-27.

FIGURE CAPTIONS

Figure 1: Clasts of Paleozoic quartzite and carbonate in conglomerate of the Colton Formation in the West Hills north of Mills Gap.

Figure 2: Oncolitic limestone in North Horn or Flagstaff strata on Skinner Peaks. (Photo by S.R. Mattox)

Figure 3: Outcrop of epiclastic conglomeratic sandstone of Unit 1 of the Goldens Ranch Formation. Note the cross-bedding, pebble lenses, and typical blue-gray color. Hammer for scale in center of photo. Photo taken in the Painted Rocks area. (Photo by S.R. Mattox)

APPENDIX 1

#### SKINNER PEAKS SECTION

add other weasured ins

This section was measured on a southwest traverse beginning on the 5700 ft contour, just south of the jeep trail in the SE 1/4 of section 15, T. 16 S., R. 1 W.; strata dip approximately 30 degrees SW.

UNIT #	UNIT	CUMULATIVE	
(SAMPLE3)	THICKNESS	THICKNESS	DESCRIPTION
13	17.0	745.0	Sandy limestone, grayish-
			yellow (5Y 8/4); slope-
			forming.
12	15.0	728.0	Calcareous sandstone, pinkish-
			gray (5YR 8/1), weathered and
			fresh; massive, ledge-
			forming;sand is 80% quartz,
			subangular to subrounded,
			moderately-sorted.
11	95.0	713.0	Sandy limestone, variable
			color; weathers into plates;
			sand is medium-grained,
			subrounded quartz.

GREEN RIVER FORMATION

#### FLAGSTAFF LIMESTONE OR NORTH HORN FORMATION

10	50.0	618.0	Interbedded pebble
			conglomerate and sandstone
			lenses; sandstone contains
			algal mat pieces (up to 5
			inches) and oncolites;
			composed of medium-grained,
			well-sorted, subangular to
			subrounded quartz;
			conglomerate clasts are 50%
			quartzite (rounded tan and
			purple from the Cambrian
			Tintic Quartzite, and the
			Precambrian Mutual Formation)
			and 50% carbonate (Paleozoic).
9	60.0	568.0	Sandy limestone and sandstone,
			<pre>pale-reddish-brown (10R 5/4);</pre>
			forms a slope with local
			ledges; sand is medium-grained
			quartz.
8	81.0	508.0	Oncolitic limestone,
			yellowish-gray (5Y 7/2);
			cliff-forming; oncolites up to
			3 inches in diameter.
7	15.0	427.0	Covered slope.
6	10.0	412.0	Limestone, finely-crystalline,

light-gray (N7); massive, ledge-forming.

#### FLAGSTAFF LIMESTONE OR NORTH HORN FORMATION

NORTH HORN FORMATION (?)

5	55.0	402.0	Conglomerate interbedded with
			sandstone; cliff and ledge-
			forming; sandstone is light-
	Sec.		gray (N7); composed of medium-
			grained, subangular to
			subrounded, well-sorted
			<pre>quartz; locally cross-bedded;</pre>
			conglomerate is clast-
			supported; 80% of the clasts
		-	are subangular to subrounded
			cobbles composed of Paleozoic
			carbonates (75%) and
			Precambrian/Cambrian quartzite
			(25%); matrix is medium-
			grained, well-sorted, rounded
			quartz sand.
4	220.0	347.0	Conglomerate; cliff and ledge-
			forming; clasts are subangular
			to subrounded pebbles,

cobbles, and boulders of purple and tan quartzite derived from the Precambrian Mutual Formation and Cambrian Tintic Quartzite respectively; matrix is coarse-grained quartz sand; unit is gray at base and changes to red upsection.

127.0 Slope covered with rubble of quartzite boulders and cobbles; derived from the conglomerate that is up-slope.

#### NORTH HORN FORMATION (?)

90.0

3

ARAPIEN SHALE

2	2.0	37.0	Limestone, finely-crystalline,
			grayish-green (10GY 5/2);
			ledge-forming; separated from
			unit 3 by a fault.
l	35.0	35.0	Calcareous mudstone, grayish-
			green (10GY 5/2).



<u>Figure 1:</u> Clasts of Paleozoic quartzite and carbonate in conglomerate of the Colton Formationin the West Hills north of Mills Gap.

Scale



<u>Figure 2:</u> Oncolitic limestone in North Horn or Flagstaff strata on Skinner Peaks. (Photo by S. R. Mattox.)



Figure 3: Outcrop of epiclastic conglomeratic sandstone of Unit I of the Goldens Ranch Formation. Note the cross-bedding, pebble lenses, and typical blue-gray color. Hammer for scale in center of photo. Photo taken in the Painted Rocks area. (Photo by S. R. Mattox.)

Scales

For Plate 2

#### DESCRIPTION OF MAP UNITS

- Qal Alluvium Clay- to boulder sized material; locally derived; occurs along most drainages.
- Qc Colluvium Steeply-sloping, cone-shaped deposits; material is unconsolidated, very angular, very poorly-sorted; color and composition reflect the formation from which the deposits were derived.
- Qls Landslide deposits Angular, poorly-sorted blocks of carbonate and sandstone in a mudstone matrix; material was derived from the Green River Formation.
- Qacf1 Younger coalescing alluvial fans Small alluvial fans Qacf1 located north of Little Salt Creek Canyon; composed of angular, pebble-sized fragments of Arapien Shale.
- Qacf2 Older coalescing alluvial fans Reddish-brown to Qafz yellowish-gray, unconsolidated, poorly-sorted clay, sand, pebbles, cobbles, and boulders; deposits are massive to crudely bedded; clasts are composed of quartzite, limestone, sandstone, and volcanic rocks.
  - Qdf Fine-grained deltaic sediments Light brown, unconsolidated, coarse- to fine-grained sand, silt, and mud

deposited by Lake Bonneville; deposits are finely laminated and cross-laminated; soft-sediment deformation structures and ripple cross-lamination are common near the base of the exposed section.

Qaf

Solitary alluvial fan - Solitary alluvial fan located in the NW corner of the quadrangle; composed of debris from the Flagstaff Formation; very dissected and faulted.

QTaf

- Old alluvial fans Poorly-sorted sand, pebbles, cobbles, and boulders; forms distinctive yellow caps in the hills north of Skinner Peaks.
- QTap Pediment alluvium Poorly sorted sand, pebbles, cobbles, and boulders; also contains red pebbly sandstone and sandy limestone; alluvium occurs as dissected caps in the South Hills.
- Tvgu Goldens Ranch Formation (undifferentiated)
- Tvg5 Unit V of the Goldens Ranch Formation Equals the Hall Canyon Conglomerate of Meibos (1983); blue-gray epiclastic conglomerate and conglomeratic sandstone; contains clasts of Unit IV.

Tvg4 Unit IV of the Goldens Ranch Formation - Orange or tan

vitric lithic tuff; contains flattened pumice up to six inches in length; weathers to vertical cliff that are commonly cavernous.

- Tvg3 Unit III of the Goldens Ranch Formation Coarse-grained red or gray epiclastic sandstone that contains cross-bedding and channels; composed of approximately 60% bipyramidal quartz crystals; forms resistant ledges.
- Tvg2 Unit II of the Goldens Ranch Formation Pink crystal vitric tuff containing biotite, bipyramidal quartz, sanidine, and pumice; weathers to form slopes.
- Tvgl Unit I of the Goldens Ranch Formation Blue-gray or green epiclastic conglomerate and conglomeratic sandstone; forms cliff and ledges that display cross-bedding and channels.
- Ti Igneous Intrusions Intrusions of hornblende monzonite porphyry; less than 30 feet in width, weather to a grus-like talus.
- Tgr Green River Formation Interbedded grayish-yellow to brown mudstone, limestone, sandstone, and conglomeratic sandstone; limestone is commonly fossiliferous or oolitic; a conspicuous bed of stromatolitic limestone occurs in the bottom part of the section; sandstone near top of section

contains vertebrate fossils.

- TKu Cretaceous and Tertiary strat (undifferentiated) Includes Tc (Colton Formation), Tf (Flagstaff Formation), and TKnh (North Horn Formation).
- Tc Colton Formation Reddish-brown mudstone, sandstone, and conglomerate; conglomerate is clast-supported, and moderately-sorted; clasts are composed of Precambrian quartzite and Paleozoic carbonate; thin beds of limestone occur locally throughout the section.
- Tf Flagstaff Formation Grayish-yellow to pale reddish-orange calcareous mudstone, sandstone, sandy limestone, limestone, and conglomerate.
- TKnh North Horn Formation Red to gray, poorly-sorted cliff and ledge-forming conglomerate; clasts are composed of quartzite and carbonate that was derived from a variety of Precambrian and Paleozoic formations. Shown only in cross-sections.
- KJu Jurassic and Cretaceous strata (undifferentiated) Includes Kpr (Price River Formation), Ki (Indianola Group), Kcm (Cedar Mountain Formation), and Jtg (Twist Gulch Formation). These units are shown only in cross-sections.

- Ja Arapien Shale Grayish-green thinly-bedded limestone, micrite, calcareous siltstone, rippled sandstone, and grayish-green or red mudstone; pods of gypsum occur locally throughout the section.
- Jtc Twin Creek Formation Shown only in cross-sections.

#### Plante 2

Form	ation	Map Symbol	Thickness	Lithology
Surficial De	Surficial Deposits		0-300	
Goldens Ranch	Unit V	Tvg₅	400 <i>- 700</i>	0;0;0; //// ©:0;0;
Formation	Unit IV	Tvay	70-100	XIII
TOTALCION	Unit III	Tuga	50-90	
	UNITI	Tvg,	40-70	·X - · X · / X · / - · X
	Unit I	Tvg,	100-500	
	Intrusive	Ti	20	
Green Riv Formati	Green River Formation		1000-1500	
Formati	N N N	Te	100 - 300	
Flagst Formati	Flagstaff Formation		100-550	
North H Formatio	orn N	TKnh	300-400	
Arapi Shale	en L	Ja	400-3000	

For Plate 2

#### MAP SYMBOLS

CONTACT

-----

Dashed where inferred; dotted where concealed

FAULT hed where inferred, dotted where conce

Dashed where inferred, dotted where concealed; bar and ball on downthrown side

> -¢-Test well

Tie-line (connects areas of like lithology)



Open -pit gypsum mine

STRIKE and DIP of BEDS

Inclined

⊕ Horizontal

-----

Vertical

For Plate 2



#### **CORRELATION OF MAP UNITS**

\* ON Cross-sections only